

FURTHER EVIDENCE THAT 1RXS J170849.0–400910 IS AN ANOMALOUS X–RAY PULSAR

G.L. ISRAEL^{1,2}, S. COVINO³, L. STELLA^{1,2}, S. CAMPANA^{3,2}, S. MEREGHETTI^{4,5}

1. Osservatorio Astronomico di Roma, Via Frascati 33, I-00040 Monteporzio Catone (Roma), Italy; Gianluca.Israel and Luigi.Stella@oar.mporzio.astro.it

2. Affiliated to the International Center for Relativistic Astrophysics (I.C.R.A.)

3. Osservatorio Astronomico di Brera, Via Bianchi 46, I-23807 Merate (Lc), Italy; covino and campana@antares.merate.mi.astro.it

4. Istituto di Fisica Cosmica “G.P.S. Occhialini” del C.N.R., Via Bassini 15, I-20133 Milano, Italy; sandro@ifctr.mi.cnr.it

Received: 25 February 1999; accepted: 15 April 1999

ABSTRACT

We report the results of two *ROSAT* HRI observations of the recently discovered ~ 11 s X–ray pulsar 1RXS J170849.0–400910. A refined position with a smaller error radius ($10''$ uncertainty) and a new spin period measurement were obtained. These results allowed to derive a period derivative of $\sim 7 \times 10^{-4}$ s yr⁻¹ and to perform a photometric and spectroscopic study of the possible optical counterparts of the source. The limits derived from the optical to X–ray flux ratio exclude the presence of a massive OB companion.

These findings, together with the nearly constant X–ray flux across observations spanning three years, strongly support the inclusion of this ~ 11 s pulsar in the class of Anomalous X–ray Pulsars (AXPs).

Subject headings: stars: neutron — pulsar: individual (1RXS J170849.0–400910) — pulsars: general — X–ray: stars

1. INTRODUCTION

The relatively bright X–ray source 1RXS J170849.0–400910 was originally discovered in the *ROSAT* All Sky Survey (RASS, Voges et al. 1998), but it did not attract much attention until ~ 11 s periodic pulsations were found during an ASCA observation that was part of the Galactic Plane Survey Project (Sugizaki et al. 1997). Based on the spin period value and the soft X–ray spectrum Sugizaki et al. (1997) suggested that this optically unidentified source could be a new member of the small group of AXPs. AXPs are a group of X–ray pulsars (Mereghetti & Stella 1995; van Paradijs et al. 1995) that share several peculiar properties: a narrow interval of pulse periods (6–12 s), very soft X–ray spectra (a steep power–law with photon index ~ 3 –4 often coupled to an additional blackbody–like component with $kT \sim 0.4$ – 0.6 keV), lower X–ray luminosities ($\lesssim 10^{35}$ erg s⁻¹) than those of persistent High Mass X–ray Binaries (HMXBs), a narrow spatial distribution in the Galactic plane, absence of large flux variations on timescales from months to years, spin–down trends at relatively stable rates (2 – 100×10^{-5} s yr⁻¹), and absence of massive companion stars (see Mereghetti et al. 1998 for a recent review). Whether AXPs are isolated neutron stars or members of binary systems remains to be determined; similarly it is unclear whether they are powered by accretion or by magnetic energy, as recently suggested by the analogy with Soft γ –ray Repeaters (SGRs; Kouveliotou et al. 1998, Mereghetti 1999).

In this Letter we report on the data analysis of public *ROSAT* HRI observations of 1RXS J170849.0–400910 and on a study of its possible optical counterparts. Our results strongly support the suggestion that 1RXS J170849.0–400910 is a member of the AXP group.

2. ROSAT HRI OBSERVATION

The field of 1RXS J170849.0–400910 was observed twice by the *ROSAT* High Resolution Imager (0.1–2.4 keV): on 1994 March 8 (ROR 900595) for a total exposure time of 1120 s and

on 1997 March 24–26 (ROR 180188) for a total exposure time of 11175 s. The source was detected at a level of 0.13 ± 0.01 and 0.130 ± 0.003 counts s⁻¹ in the 1994 and 1997 observations, respectively (90% uncertainties are used through out this letter).

We analyzed the two HRI images using both a sliding cell and a wavelet transform–based detection algorithm (Lazzati et al. 1999; Campana et al. 1999). In the longer observation (1997) the source was detected at R.A. = $17^h 08^m 47^s.24$ and DEC = $-40^\circ 08' 50''.7$ (J2000), with a statistical error of only $0.15''$ (1σ). However, due to the the uncertainty in the satellite boresight, an overall error radius of $\sim 10''$ must be considered. The new position is located some $35''$ and $25''$ North of the ASCA position ($30''$ radius; Sugizaki et al. 1997) and PSPC RASS position ($9''$ uncertainty). The 1993 *ROSAT* HRI observation was analyzed in the same way and provided a slightly different (but consistent) position, R.A. = $17^h 08^m 46^s.52$ and DEC = $-40^\circ 08' 52''.5$ (J2000; $\sim 10''$ radius uncertainty; see also Fig. 3).

The *ROSAT* HRI has no intrinsic spectral resolution, therefore in order to obtain the source flux from the observed count rate we assumed the spectral model derived with ASCA (Sugizaki et al. 1997), i.e. a power–law with photon index $\Gamma = 3.45$ and $N_H = 1.8 \times 10^{22}$ cm⁻². In this way we determined a flux at the Earth of 1.1×10^{-11} erg cm⁻² s⁻¹ in the 0.1–2.4 keV energy range. For a distance of 10 kpc, this corresponds to a source unabsorbed luminosity of $L_X \sim 1.2 \times 10^{36}$ erg s⁻¹ in the 0.8–10 keV range.

The event list of 1RXS J170849.0–400910 was extracted from a circle of $\sim 40''$ radius (corresponding to an encircled energy of $\sim 90\%$) around the X–ray position of the 1997 observation. The photon arrival times were corrected to the barycentre of the solar system and a background subtracted light curve was accumulated in 1 s bins. A power spectrum was calculated over the entire observation duration. A highly significant peak ($\sim 8\sigma$) was found at a frequency of 0.09092712 Hz, corresponding to a period of 10.998 s. To refine the period

⁵The results reported in this Letter are partially based on observations carried out at ESO, La Silla, Chile (61.D–0513).

determination and reduce its uncertainty we adopted a phase fitting technique. The best pulse period was determined to be $P=10.99802\pm0.00005$ s. The shape of the modulation is somewhat asymmetric with a pulsed fraction of $\sim 38\pm 4\%$ (Fig. 1). We note that such pulse shape is similar to that reported by Sugizaki et al. in the 0.8–2 keV range (1997; see their Fig. 3). Owing to poor statistics the periodicity was not detected during the 1994 observation (assuming a 3σ detection threshold).




FIG. 1.— The 1997 March 24–26 ROSAT HRI light curve folded at the best period $P=10.99802$ s.




FIG. 2.— Color-Magnitude Diagram for the objects in the field of 1RXS J170849.0–400910. The Main Sequence lines for different values of the distance and reddening are also shown (700 pc, $A_V=1.6$; 3 kpc, $A_V=6.2$). The straight line indicates the direction of the reddening and corresponds to a value of $A_V=2$. The squares mark the objects within the error circles.

By comparing our period value with the one measured with ASCA on 1996 September 3 (Sugizaki et al. 1997) a period derivative $\dot{P} = (7\pm 2)\times 10^{-4}$ s yr $^{-1}$ was inferred.

3. OPTICAL IMAGING AND SPECTROSCOPY

Photometry for each stellar object in the images was derived by means of DAOPHOT II (Stetson 1987), and a Color-Magnitude Diagram (CMD) was then computed (see Fig. 2) for all stars within a region of $4'\times 5'$ around the X-ray position of 1RXS J170849.0–400910. For comparison Fig. 2 also shows the main sequence for two representative distances: 700 pc (with $A_V=1.6$) and 3 kpc ($A_V=6.2$). Note that 1RXS J170849.0–400910 is located in the Galactic plane ($l''=346.48$, $b''=0.03$) and that along this direction spiral arms are located at a distance of ~ 1 kpc, ~ 3 kpc and ~ 4.5 kpc (Taylor & Cordes 1993). The R image of the field of 1RXS J170849.0–400910 is shown in Fig. 3. The new X-ray position uncertainty regions obtained with the two HRI observations are shown ($10''$ radius). Table 1 gives the positions, R magnitudes and the colors of the objects within (or close to) the error circles.

The spectrum of the brightest object within the error circle (candidate A, $R = 13.4$; $V-R = 1.48$) is shown in Fig. 4. The object was classified as a B0V spectral-type star with an absorption $A_V \sim 6.2$. We found no evidence for any emission-lines in its spectrum.




FIG. 3.— R image of the field of 1RXS J170849.0–400910. The new X-ray position uncertainty regions ($10''$ radius) obtained with the 1994 and 1997 HRI observations are also shown. North is top, east is left.

4. DISCUSSION

The great majority of accreting X-ray pulsars are found in two distinct classes of massive binaries: systems with OB type supergiant companions and systems with Be stars. The latter are often characterized by transient activity.

We found that star A, the brightest candidate counterpart in the new error circles, is a main sequence B0 star, but the properties of its spectrum make an association with 1RXS J170849.0–400910 very unlikely. The absence of emission-lines corresponding to the Balmer series, HeI and FeII strongly suggest that star A is not in a binary system with a compact object. Although the optical emission-lines in Be/neutron star systems may occasionally disappear during the quiescence state, such a behaviour has been so far observed only in transient sources which show also outbursts and pronounced spin-up/down rate




FIG. 4.— 4300–6800 Å low-resolution flux-calibrated spectrum of the reddened B0V spectral-type star A. I.S. stands for inter-stellar absorption.

episodes. The low value of the inferred spin-down period derivative over a 6 month interval and the constant X-ray flux value over a timescale of 3 years (see below) rule against this possibility. Furthermore, the soft X-ray spectrum of 1RXS J170849.0–400910 is much softer than that observed in all X-ray pulsars with massive companions (i.e. a hard power law with photon index ~ 1 that steepens only above the cut-off energy of ~ 10 – 30 keV).

TABLE 1

POSITIONS, R MAGNITUDES AND COLORS OF THE STARS WITHIN (OR CLOSE TO) THE X-RAY ERROR CIRCLES OF 1RXS J170849.0–400910

Star	R.A. (J2000) ^a (hh mm ss)	DEC (J2000) ^a (° ' ")	R ^b mag	V–R ^b mag
A	17 08 46.2	–40 08 48	13.41	1.48
B	17 08 47.8	–40 08 40	16.46	0.85
C	17 08 46.8	–40 08 40	17.36	1.48
D	17 08 47.1	–40 08 36	18.24	1.50
E	17 08 47.6	–40 08 46	18.26	1.27
F	17 08 46.1	–40 08 43	18.36	1.38
G	17 08 47.0	–40 08 38	18.60	1.50
H	17 08 46.9	–40 09 05	18.74	1.25
I	17 08 46.4	–40 08 53	18.86	1.61
J	17 08 46.8	–40 08 50	18.93	1.36
K	17 08 47.8	–40 08 52	19.00	1.38
L	17 08 47.4	–40 08 53	19.78	1.58

^a— Uncertainty $\sim 1''$.

^b— Absolute uncertainty ~ 0.2 mag. Relative uncertainty ~ 0.02 mag.

The second brightest candidate to be considered is star B. Its position in the CMD of Fig. 2 is consistent with a relatively closely K type star.

All remaining candidates are fainter than $R \sim 17$ implying X-ray to optical flux ratios higher than those typically observed in High Mass X-ray Binaries. This is also indicated by their position in the CMD shown in Fig. 2: the objects in the error circles are too blue and too faint to be compatible with early type stars. Note also that our images are deep enough to exclude very reddened and distant O and B stars. The absorption column density for 1RXS J170849.0–400910 is $\sim 1.5 \times 10^{22} \text{ cm}^{-2}$

(Sugizaki et al. 1997), corresponding to $E(B-V) \sim 2.6$ (Bohlin et al. 1978). Applying the standard interstellar extinction law (Fitzpatrick 1998) this translates to an absorption in the R band of $A_R \sim 6$. A B0 star located at a distance of $5 \div 10$ kpc would be observed with a R magnitude $\sim 15.5 \div 17$ and $V-R \sim 2$, i.e. it should have been easily detected in our optical images.

Our optical data cannot exclude that 1RXS J170849.0–400910 is part of a Low Mass X-ray Binary (LMXB). The faint stars in the error box are all compatible with relatively late spectral-type stars. For example star I is consistent with a A5–7V at a distance of 3 kpc, while star H is consistent with a late K or an early M at a distance of 700 pc. However, none of these objects has the characteristic blue color generally observed in the counterparts of luminous LMXBs, in which the optical emission is dominated by the accretion disk (van Paradijs 1998).

The lack of a massive companion star and the inferred spin-down at a rate of $\sim 7.5 \times 10^{-4} \text{ s yr}^{-1}$ strongly support the inclusion of 1RXS J170849.0–400910 in the group of AXPs, as was previously suggested by Sugizaki et al. (1997) based only on the soft X-ray spectrum and 11 s period.

1RXS J170849.0–400910 was detected by the *ROSAT* HRI at the same flux level in both the 1994 and 1997 observations. The *ROSAT* flux extrapolated to the 0.8–10 keV range ($2.3 \times 10^{-11} \text{ erg cm}^{-2} \text{ s}^{-1}$; for the power-law best fit) is about a factor of 2 lower than that observed with *ASCA* ($4.3 \times 10^{-11} \text{ erg cm}^{-2} \text{ s}^{-1}$). A soft blackbody component, accounting for 20–50% of the total flux has been seen in the spectrum of three well studied AXPs (4U 0142+61; White et al. 1996; Israel et al. 1999, 2E 2259+587; Corbet et al. 1995; Parmar et al. 1998, and 1E 1048.1–5937; Corbet & Mihara 1997; Oosterbroek et al. 1998). The presence of a similar component in 1RXS J170849.0–400910 was found compatible with the *ASCA* data (although not formally required by the fit, Sugizaki et al. 1997). Assuming a double component spectral model for 1RXS J170849.0–400910, at least part of the flux difference between the *ASCA* and *ROSAT* observations might be ascribed to the more complex spectral slope than assumed. We conclude that the modest luminosity variations across observations spanning three years is another analogy of 1RXS J170849.0–400910 with the AXPs class. 1RXS J170849.0–400910, like other AXPs, displays also a remarkable stability of the pulsed fraction and pulse shape.

One of the main problems in understanding the nature of AXPs is to establish whether they are isolated neutron stars or members of binary systems with very low mass companions. Both deep optical and infrared searches for counterparts

and searches for Doppler orbital modulations in the pulse frequency can provide valuable information in this respect. In the best studied AXPs, the limits derived from such studies exclude most classes of companion stars, but helium stars with $M \leq 0.8 M_{\odot}$ and white dwarves (Mereghetti, Israel & Stella 1998; Wilson et al. 1999).

The possibility that 1RXS J170849.0–400910 has a white dwarf companion star is compatible with our optical observations: the expected R magnitude at a distance of 700 pc would be ≥ 21 , well below the limiting magnitude of our images. Similar results apply to a helium main sequence star. We also note that although the absorption column inferred from the ASCA data would place 1RXS J170849.0–400910 quite far from us

(10 kpc), it is possible that part of the absorption be intrinsic. Also the possibility that 1RXS J170849.0–400910 is a “magnetar”, an isolated strongly magnetic neutron star, possibly related to Soft γ –ray Repeaters remains viable. More period measurements are clearly required to look for “glitches” as foreseen in the “magnetar” scenario. In conclusion, based on the X–ray and optical findings presented here, we confirm that 1RXS J170849.0–400910 is a new member of the AXPs group.

GLI and SC thanks V.F. Polcaro for his kind help and useful discussions. This work was partially supported through ASI grants.

REFERENCES

- Bohlin, R.C., Savage, B.D., and Drake, J.F., 1978, *ApJ*, 224, 132
 Campana, S., Lazzati, D., Panzera, R., et al., 1999, *ApJ*, submitted
 Corbet, R.H.D., Smale, A.P., Ozaki, M., et al., 1995, *ApJ*, 443, 786
 Corbet, R.H.D., and Mihara, T., 1997, *ApJ*, 475, L127
 Fitzpatrick, E.L., 1998, *astro-ph/9809387*
 Israel, G.L., Oosterbroek, T., Angelini, L., et al., 1999, *A&A*, in press (*astro-ph/9904103*)
 Lazzati, D., Campana, S., Rosati, P., et al., 1999, *ApJ*, submitted
 Kouveliotou, C., Dieters, S., Strohmayer, T., et al., 1998, *Nature*, 393, 235
 Mereghetti, S., 1999, in “Relationship between Neutron Stars and Supernova Remnants”, *Mem.S.A.It.*, in press (<http://www.arcetri.astro.it/~elba98/index.html>)
 Mereghetti, S., and Stella, L., 1995, *ApJ*, 442, L17
 Mereghetti, S., Israel, G.L., and Stella, L., 1998, *MNRAS* 296, 689
 Mereghetti, S., Stella, L., and Israel, G.L., 1998, in “The Active X–ray Sky: Results from BeppoSAX and RossiXTE”, *Nuclear Physics B Proceedings Supplements*, Scarsi, L., Bradt H., Giommi, P., and Fiore, F. (eds), Elsevier Science B.V. (Amsterdam), p. 253
 Oosterbroek, T., Parmar, A.N., Mereghetti, S., et al., 1998, *A&A*, 334, 925
 Parmar, A.N., Oosterbroek, T., Favata, F., et al., 1998, *A&A*, 330, 175
 Stetson, P.B., 1987, *PASP*, 99, 191
 Sugizaki, M., Nagase, F., Torii, K., et al., 1997, *PASJ*, 49, L25
 Taylor, J.H., and Cordes, J.M., 1993, *ApJ*, 411, 674
 van Paradijs, J., Taam, R.E., and van den Heuvel, E.P.J., 1995, *A&A*, 299, L41
 van Paradijs, J., 1998, in “The Many Faces of the Neutron Stars”, Buccheri, R., van Paradijs, J., Alpar, M.A. (eds), Kluwer Academic Publishers (The Netherlands), p.279
 Voges, W., Aschenbach, B., Boller, T.H., et al., 1998, in “New Horizons from Multi–Wavelength Sky Surveys”, 179th Symposium of the International Astronomical Union, McLean, B.J., Golombek, D.A., Hayes, J.J.E., and Payne, H.E. (eds), Kluwer Academic Publishers, p.33
 White, N.E., Angelini, L., Ebisawa, K., et al., 1996, *ApJ*, 463, L83
 Wilson, C.A., Dieters, S., Finger, M., et al., 1999, *ApJ*, 513, 464